

Rabbit anti-DDDDK tag antibody, clone SQab1880 (monoclonal)

Clone no. SQab1880

MONOSAN

| | |
|---------------------------|---|
| Product name | Rabbit anti-DDDDK tag antibody, clone SQab1880 (monoclonal) |
| Host | Rabbit |
| Applications | FC, ICC/IF, IP, WB |
| Species reactivity | Other |
| Conjugate | - |
| Immunogen | KLH-conjugated DYKDDDDK. |
| Isotype | - |
| Clonality | Monoclonal |
| Clone number | SQab1880 |
| Size | 100 ul |
| Concentration | n/a |
| Format | Purification with Protein A. |
| Storage buffer | PBS, 0.01% Sodium azide, 40% Glycerol and 0.05% BSA. |
| Storage until expiry date | -20°C |

FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES

Rabbit anti-DDDDK tag antibody, clone SQab1880 (monoclonal)


Clone no. SQab1880

MONOSAN


Additional info

Application note: * The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist. Storage instruction: For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot and store at -20°C. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use. Background:

Images



Immunofluorescence: 293 cells transfected with C-terminal DYKDDDDK tagged gene. Cells were fixed with 4% paraformaldehyde for 30 min at RT, permeabilized with 0.1% Triton X-100 for 10 min at RT then blocked with 10% goat serum for 30 min at RT. Cells were stained with DAPI (left)



Western blot: Testing of anti-DDDDK tag antibody [SQab1880]. 3 µg of 293 cell lysate without any transfection (at 1:2000, left) or transfected with N-terminal DYKDDDDK tagged gene (at 1:10000, right).

References

1. -
2. -
3. -
4. -
5. -

FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES