

Rabbit anti-vWF antibody, clone SQab19164 (monoclonal)

Clone no. SQab19164

MONOSAN

---

|                           |  |
|---------------------------|--|
| Product name              | Rabbit anti-vWF antibody, clone SQab19164 (monoclonal)                       |
| Host                      | Rabbit   |
| Applications              | IHC-P  |
| Species reactivity        | Human  |
| Conjugate                 | -  |
| Immunogen                 | Synthetic peptide within aa. 1150-1250 of Human von Willebrand factor (vWF). |
| Isotype                   | -  |
| Clonality                 | Monoclonal   |
| Clone number              | SQab19164  |
| Size                      | 100 ul   |
| Concentration             | n/a  |
| Format                    | Purification with Protein A.   |
| Storage buffer            | PBS, 0.01% Sodium azide, 40% Glycerol and 0.05% BSA.                         |
| Storage until expiry date | -20°C  |

FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES

Rabbit anti-vWF antibody, clone SQab19164 (monoclonal)

Clone no. SQab19164


MONOSAN

---

**Additional info**

Application note: IHC-P: Antigen Retrieval: Heat mediation was performed in Tris/EDTA buffer (pH 9.0), primary antibody incubate at RT (18°C-25°C) for 30 minutes.\* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist. Storage instruction: For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot and store at -20°C. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use. Background: The glycoprotein encoded by this gene functions as both an antihemophilic factor carrier and a platelet-vessel wall mediator in the blood coagulation system. It is crucial to the hemostasis process. Mutations in this gene or deficiencies in this protein result in von Willebrand's disease. An unprocessed pseudogene has been found on chromosome 22. [provided by RefSeq, Jul 2008]

## Images



Immunohistochemistry: Formalin/PFA-fixed and paraffin-embedded Human cervix tissue stained with anti-vWF antibody [SQab19164]. Antigen Retrieval: Heat mediation was performed in Tris/EDTA buffer (pH 9.0).

-

0

## References

1. -
2. -
3. -
4. -
5. -

FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES