

Rabbit anti-ALK antibody, clone SQab20239 (monoclonal)

Clone no. SQab20239

MONOSAN

| | |
|---------------------------|--|
| Product name | Rabbit anti-ALK antibody, clone SQab20239 (monoclonal) |
| Host | Rabbit |
| Applications | IHC-P |
| Species reactivity | Human |
| Conjugate | - |
| Immunogen | Recombinant protein of Human ALK. |
| Isotype | - |
| Clonality | Monoclonal |
| Clone number | SQab20239 |
| Size | 100 ul |
| Concentration | n/a |
| Format | Purification with Protein A. |
| Storage buffer | PBS, 0.01% Sodium azide, 40% Glycerol and 0.05% BSA. |
| Storage until expiry date | -20°C |

FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES

Rabbit anti-ALK antibody, clone SQab20239 (monoclonal)


Clone no. SQab20239

MONOSAN

Additional info

Application note: IHC-P: Antigen Retrieval: Heat mediation was performed in Tris/EDTA buffer (pH 9.0), primary antibody incubate at RT (18°C - 25°C) for 30 minutes.* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist. Storage instruction: For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot and store at -20°C. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use. Background: This gene encodes a receptor tyrosine kinase, which belongs to the insulin receptor superfamily. This protein comprises an extracellular domain, an hydrophobic stretch corresponding to a single pass transmembrane region, and an intracellular kinase domain. It plays an important role in the development of the brain and exerts its effects on specific neurons in the nervous system. This gene has been found to be rearranged, mutated, or amplified in a series of tumours including anaplastic large cell lymphomas, neuroblastoma, and non-small cell lung cancer. The chromosomal rearrangements are the most common genetic alterations in this gene, which result in creation of multiple fusion genes in tumourigenesis, including ALK (chromosome 2)/EML4 (chromosome 2), ALK/RANBP2 (chromosome 2), ALK/ATIC (chromosome 2), ALK/TFG (chromosome 3), ALK/NPM1 (chromosome 5), ALK/SQSTM1 (chromosome 5), ALK/KIF5B (chromosome 10), ALK/CLTC (chromosome 17), ALK/TPM4 (chromosome 19), and ALK/MSN (chromosome X). [provided by RefSeq, Jan 2011]

Images



Immunohistochemistry: Formalin/PFA-fixed and paraffin-embedded Human non-small-cell lung carcinoma tissue. Antigen Retrieval: Heat mediation was performed in Tris/EDTA buffer (pH 9.0). The tissue section was stained with anti-ALK antibody [SQab20239].

-

0

References

1. -
2. -
3. -
4. -
5. -

FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES