

MONOCLONAL ANTIBODY

# PE labeled Anti-Human Podocalyxin/PCLP1

Code No.	Clone	Subclass	Quantity
M084-5	53D11	Mouse IgG2a	1 mL (50 tests)

**BACKGROUND:** Recent studies with avian embryos and murine embryonic stem cells have suggested that hematopoietic cells are derived from hemangioblasts, the common precursors of hematopoietic and endothelial cells. Hara *et al.* molecularly cloned podocalyxin-like protein 1 (PCLP1) as a novel surface marker for endothelial-like cells in the AGM (aorta-gonad-mesonephros) region of mouse embryos, where long-term repopulating hematopoietic stem cells (LTR-HSCs) are known to arise. PCLP1<sup>+</sup>CD45<sup>-</sup> cells in the AGM region incorporated acetylated low-density lipoprotein and produced both hematopoietic and endothelial cells when cocultured with OP9 stromal cells. Moreover, multiple lineages of hematopoietic cells were generated in vivo when PCLP1<sup>+</sup>CD45<sup>-</sup> cells were injected into neonatal liver of busulfan-treated mice. Today it is reported that the PCLP1 is identical with the Podocalyxin.

**SOURCE:** This antibody was purified from mouse ascites fluid using protein A agarose. This hybridoma (53D11) was established by fusion of mouse myeloma cell P3U1 with Balb/c mouse splenocyte immunized with CHO cell expressing full-length human Podocalyxin/PCLP1.

**FORMULATION:** 50 tests in 1 mL volume of PBS containing 1% BSA and 0.09% NaN<sub>3</sub>.

\*Azide may react with copper or lead in plumbing system to form explosive metal azides. Therefore, always flush plenty of water when disposing materials containing azide into drain.

**STORAGE:** This antibody solution is stable for one year from the date of purchase when stored at 4°C.

**REACTIVITY:** This antibody reacts with human Podocalyxin/PCLP1 on Flow cytometry.

**APPLICATION:**

Flow cytometry; 20 µL (ready for use)

\*Please refer to the data sheet (MBL; code no. M084-3) for other applications.

Detailed procedure is provided in the following **PROTOCOL.**

**INTENDED USE:**

For Research Use Only. Not for use in diagnostic procedures.

**SPECIES CROSS REACTIVITY:**


Species	Human	Mouse	Rat
Cell	HUVEC	Not Tested	Not Tested
Reactivity on FCM	+		

**REFERENCES:**

- 1) Doyonnas, R., *et al.*, *Blood* **105**, 4170-4178 (2005)
- 2) Minegishi, N., *et al.*, *Blood* **102**, 896-905 (2003)
- 3) Schopperle, M.W., *et al.*, *Biochem. Biophys. Res. Commun.* **300**, 285-290 (2003)
- 4) Minehata, K., *et al.*, *Blood* **99**, 2360-2368 (2002)
- 5) Doyonnas, R., *et al.*, *J. Exp. Med.* **194**, 13-27 (2001)
- 6) Hara, T., *et al.*, *Immunity* **11**, 567-578 (1999)
- 7) Kershaw, B. D., *et al.*, *J. Biol. Chem.* **272**, 15708-15714 (1997)

**RELATED PRODUCTS:**

- M084-3 Anti-Human Podocalyxin/PCLP1 (53D11)
- M084-4 FITC labeled anti-Human Podocalyxin/PCLP1 (53D11)
- M085-3 Anti-Human Podocalyxin/PCLP1 (4H11)
- D072-3 Anti-Mouse Podocalyxin/PCLP1 (10B9)
- D072-4 FITC labeled anti-Mouse Podocalyxin/PCLP1 (10B9)
- D072-5 PE labeled anti-Mouse Podocalyxin/PCLP1 (10B9)
- D072-6 Biotin labeled anti-Mouse Podocalyxin/PCLP1 (10B9)



**Flow cytometric analysis of PCLP1 expression on HUVEC (left) and CHO (right).** Open histogram indicates the reaction of isotypic control to the cells. Shaded histograms indicate the reaction of M084-5 to the cells.

**PROTOCOL:**

**Flow cytometric analysis for adherent cells**

We usually use Fisher tubes or equivalents as reaction tubes for all steps after 2).

- 1) Detach the cells from culture dish by using cell dissociation buffer (Invitrogen; code no. 13151-014).
- 2) Wash the cells 3 times with washing buffer [PBS

- containing 2% fetal calf serum (FCS) and 0.1% NaN<sub>3</sub>].
- 3) Resuspend the cells with washing buffer (5x10<sup>6</sup> cells/mL).
  - 4) Add 50 µL of the cell suspension into each tube, and centrifuge at 500 x g for 1 minute at room temperature. Remove supernatant by careful aspiration.
  - 5) Add 20 µL of Clear Back (human Fc receptor blocking reagent, MBL; code no. MTG-001) to the cell pellet after tapping. Mix well and incubate for 5 minutes at room temperature.
  - 6) Add 20 µL of the PE labeled anti-Human podocalyxin/PCLP1 monoclonal antibody (53D11). Mix well and incubate for 30 minutes at room temperature.
  - 7) Add 1 mL of the washing buffer followed by centrifugation at 500 x g for 1 minute at room temperature. Remove supernatant by careful aspiration.
  - 8) Resuspend the cells with 500 µL of the washing buffer and analyze by a flow cytometer.

(Positive control for Flow cytometry; HUVEC)