

POLYCLONAL ANTIBODY

# Anti-Human IL-33

<b>Code No.</b>	<b>Quantity</b>	<b>Form</b>
PM033	100 µL	Affinity Purified

**BACKGROUND:** Interleukin-1 (IL-1) family, such as IL-1 $\alpha$ / $\beta$  and IL-18, have important functions in host defense, immune regulation, and inflammation. IL-33, a member of the IL-1 family, that shows to induce T helper (Th) type 2 responses by signaling through the IL-1 receptor-related protein ST2 (IL-1R4), an orphan member of the IL-1 receptor family. Similarly to IL-1 $\alpha$ / $\beta$  and IL-18, IL-33 is synthesized as a 31 kDa precursor protein has been shown to be cleaved by caspase-1 in vitro. In vivo, IL-33 induces the expression of IL-4, IL-5, and IL-13 and leads to severe pathological changes in mucosal organs. IL-33 has been originally identified as NF-HEV, which is a nuclear factor preferentially expressed in high endothelial venules. IL-33 may function as both a proinflammatory cytokine and an intracellular nuclear factor involved in transcriptional regulation.

**SOURCE:** This antibody was purified from rabbit serum using affinity column. The rabbit was immunized with the recombinant full-length human IL-33 (270 aa).

**FORMULATION:** 100 µL volume of PBS containing 50% glycerol, pH 7.2. No preservative is contained.

**STORAGE:** This antibody solution is stable for one year from the date of purchase when stored at -20°C.

**REACTIVITY:** This antibody reacts with IL-33 on Western blotting.

**APPLICATIONS:**

Western blotting; 1:200 for chemiluminescence detection system

Immunoprecipitation; Not recommended

Immunohistochemistry; 1:500 - 1:1,000

Heat treatment is necessary for paraffin embedded sections.

Autoclave; 10 minutes at 110 °C in 10 mM citrate buffer (pH 6.5)

Immunocytochemistry; Not tested

Flow cytometry; Not tested

Detailed procedure is provided in the following **PROTOCOLS**.

**INTENDED USE:**


For Research Use Only. Not for use in diagnostic procedures.

**SPECIES CROSS REACTIVITY:**

Species	Human	Mouse	Rat
Cells	Raji, HeLa, HUVEC, Lu99A	Not Tested	Not Tested
Reactivity on WB	+		

**REFERENCE:**

- 1) Schmitz, J., *et al.*, *Immunity* **23**, 479–490 (2005)



**Western blot analysis of IL-33 expression in transfectant (1), Raji (2), HeLa (3), HUVEC (4) and Lu99A (5) using PM033.**


**PROTOCOLS:**

**SDS-PAGE & Western Blotting**

- 1) Wash the 1x10<sup>7</sup> cells 3 times with PBS and suspend with 1 mL of Laemmli's sample buffer.
- 2) Boil the samples for 2 minutes and centrifuge. Load 20 µL of the sample per lane in a 1 mm thick SDS-polyacrylamide gel for electrophoresis.
- 3) Blot the protein to a polyvinylidene difluoride (PVDF) membrane at 1 mA/cm<sup>2</sup> for 1 hour in a semi-dry transfer system (Transfer Buffer: 25 mM Tris, 190 mM glycine, 20% MeOH). See the manufacture's manual for precise transfer procedure.
- 4) To reduce nonspecific binding, soak the membrane in 5% skimmed milk (in PBS, pH 7.2) for 1 hour at room temperature, or overnight at 4°C.
- 5) Incubate the membrane with primary antibody diluted with PBS, pH 7.2 containing 1% skimmed milk as suggest in the **APPLICATIONS** for 1 hour at room temperature. (The concentration of antibody will depend on condition.)

- 6) Wash the membrane with PBS-T [0.05% Tween-20 in PBS] (5 minutes x 3 times).
- 7) Incubate the membrane with the 1:10,000 HRP-conjugated anti-rabbit IgG (MBL; code no. 458) diluted with 1% skimmed milk (in PBS, pH 7.2) for 1 hour at room temperature.
- 8) Wash the membrane with PBS-T (5 minutes x 3 times).
- 9) Wipe excess buffer on the membrane, then incubate it with appropriate chemiluminescence reagent for 1 minute. Remove extra reagent from the membrane by dabbing with paper towel, and seal it in plastic wrap.
- 10) Expose to an X-ray film in a dark room for 10 minutes. Develop the film as usual. The condition for exposure and development may vary.

(Positive controls for Western blotting; transfectant, 293T, Raji, HeLa, HUVEC, Lu99A)



**Immunohistochemical detection of IL-33 on human tonsil paraffin embedded section with PM033.**

#### **Immunohistochemical staining for paraffin-embedded sections: SAB method**

- 1) Deparaffinize the sections with Xylene 3 times for 3-5 minutes each.
- 2) Wash the slides with Ethanol 3 times for 3-5 minutes each.
- 3) Wash the slides with PBS 3 times for 3-5 minutes each.
- 4) Heat treatment  
Heat treatment by Autoclave:  
Place the slides put on staining basket in 500 mL beaker with 500 mL of 10 mM citrate buffer (pH 6.5). Cover the beaker with plastic wrap, then process the slides with the autoclave for 10 minutes at 110°C. Let the slides cool down in the beaker at room temperature for about 40 minutes.
- 5) Remove the slides from the citrate buffer and cover each section with 3% H<sub>2</sub>O<sub>2</sub> for 10 minutes at room temperature to block endogenous peroxidase activity. Wash 3 times in PBS for 5 minutes each.
- 6) Remove the slides from PBS, wipe gently around each section and cover tissues with blocking buffer (1% BSA,

- 20 mM Hepes, 135 mM NaCl) for 10 minutes to block non-specific staining. Do not wash.
- 7) Tip off the blocking buffer, wipe gently around each section and cover tissues with primary antibody diluted with blocking buffer as suggest in the **APPLICATIONS**.
- 8) Incubate the sections for 1 hour at room temperature.
- 9) Wash the slides 3 times in PBS for 5 minutes each.
- 10) Wipe gently around each section and cover tissues with enzyme conjugated polymer reagent. Incubate for 1 hour at room temperature. Wash as in step 9).
- 11) Visualize by reacting for 10-20 minutes with substrate solution containing 7.5 mg DAB, 40 µL of 30% H<sub>2</sub>O<sub>2</sub> in 150 mL PBS. \*DAB is a suspect carcinogen and must be handled with care. Always wear gloves.
- 12) Wash the slides in water for 5 minutes.
- 13) Counter stain in hematoxylin for 1 minute, wash the slides 3 times in water for 5 minutes each, and then immerse the slides in PBS for 5 minutes. Dehydrate by immersing in Ethanol 3 times for 3 minutes each, followed by immersing in Xylene 3 times for 3 minutes each.
- 14) Now ready for mounting.

(Positive control for Immunohistochemistry; human tonsil)

#### **RELATED PRODUCTS:**

- D046-3 Anti-Mouse IL-18 (39-3F)
- D047-3 Anti-Mouse IL-18 (74)
- D048-3 Anti-Mouse IL-18 (93-10C)
- D048-6 Biotin labeled anti-Mouse IL-18 (93-10C)
- D043-3 Anti-Human IL-18 (25-2G)
- D044-3 Anti-Human IL-18 (125-2H)
- D045-3 Anti-Human IL-18 (159-12B)
- D045-6 Biotin labeled anti-Human IL-18 (159-12B)
- PM014 Anti-Human IL-18 (poly)
- 7620 Anti-Human IL-18 ELISA Kit
- 7625 Anti-Mouse IL-18 ELISA Kit
- B001-5 Recombinant Human IL-18
- B003-5 Recombinant Human IL-18 (without BSA)
- B002-5 Recombinant Mouse IL-18
- B004-5 Recombinant Mouse IL-18 (without BSA)
- D065-3 Anti-ST2 (HB12)
- D066-3 Anti-ST2 (FB9)
- D067-3 Anti-ST2 (2A5)
- D067-4 FITC labeled anti-ST2 (2A5)
- D067-5 PE labeled anti-ST2 (2A5)
- 7638 ST2 ELISA Kit